

Author Index

Barañao, J.L., see Pomata, P.E. (120) 83
Barbosa, M., see Villalobos, J. (120) 99
Barker, J.L., see Liu, Q.-Y. (120) 223
Barr, G.A., see Wiedenmayer, C.P. (120) 191
Bezin, L., see Groc, L. (120) 95
Bishara, O., see Thomas, A.J. (120) 181
Boylan, C.B., Kesterson, K.L., Crnko-Hoppen-

Boylan, C.B., Kesterson, K.L., Crnko-Hoppenjans, T.A., Ke, M., Rizk, T., Mooney, R.D. and Rhoades, R.W.

The cortical vibrissae representation is normal in transgenic mice lacking the 5-HT_{1B} receptor (120) 91

Brusco, A., see Ríos, H. (120) 17 Brzezinski, M.R., see Trofimova-Griffin, M.E. (120) 7

Cadet, N. and Paquin, J.

Conversion and storage of somatostatin are established before response to secretagogue stimuli in P19 neurons (120) 211

Cernuda-Cernuda, R., see Llamosas, M.M. (120) 1

Charli, J.-L., see Niquet, J. (120) 49 Cherry, S.R., see Moore, A.H. (120) 141 Chou, D.K.H., see Zhao, Z. (120) 165 Colman-Lerner, A.A., see Pomata, P.E. (120)

Cork, R.J., see Wu, H.H. (120) 105 Coulombe, M., see Liu, Q.-Y. (120) 223 Crnko-Hoppenjans, T.A., see Boylan, C.B. (120) 91

Dumm, J., see Liu, Q.-Y. (120) 223

Eliyayev, Y., see Gilad, V.H. (120) 255 Ernsberger, P., see Thomas, A.J. (120) 181 Erokwu, B.O., see Thomas, A.J. (120) 181 Eto, K., see Fukuda, T. (120) 65

Fernández-Ruiz, J., see Pérez-Rosado, A. (120) 77

Fiszman, M.L., see Pomata, P.E. (120) 83 Fleming, D.E., see Lund, T.D. (120) 261 Fosser, N.S., see Ríos, H. (120) 17 Foster, J.A., see Groc, L. (120) 95 Fox, E.A.

The previously identified r3/r5 repressor may require the cooperation of additional negative elements for rhombomere restriction of *Hoxb1* (120) 151

Fukuda, T., Kawano, H., Osumi, N., Eto, K. and Kawamura, K. Histogenesis of the cerebral cortex in rat fetuses with a mutation in the *Pax-6* gene (120) 65

García-Fernández, J.M., see Llamosas, M.M. (120) 1

Gilad, G.M., see Gilad, V.H. (120) 255 Gilad, V.H., Rabey, J.M., Eliyayev, Y. and

Gilad, G.M.
Different effects of acute neonatal stressors and long-term postnatal handling on stress-induced changes in behavior and in ornithine decarboxylase activity of adult rats (120) 255

Goodwin, G.A., see Wiedenmayer, C.P. (120) 191

Groc, L., Levine, R.A., Foster, J.A., Normile, H.J., Weissmann, D. and Bezin, L. Evidence of deprenyl-insensitive apoptosis of nigral dopamine neurons during development (120) 95

Grouselle, D., see Niquet, J. (120) 49 Guerra, M., see Niquet, J. (120) 49

Hayashi, H., see Orikasa, C. (120) 245 Hayashi, S., see Orikasa, C. (120) 245 Hongo, S., see Nishinaka, N. (120) 57 Hovda, D.A., see Moore, A.H. (120) 141 Huang, P.L., see Wu, H.H. (120) 105 Huerta, J.J., see Llamosas, M.M. (120) 1

Jaworski, D.M. and Proctor, M.D.

Developmental regulation of pituitary adenylate cyclase-activating polypeptide and PAC₁ receptor mRNA expression in the rat central nervous system (120) 27

Jaynes, C.D. and Turner, J.E.

Isolation of a retinal pigment epithelial
cell-derived fraction which promotes Müller
cell proliferation (120) 267

Johnson, F., Norstrom, E. and Soderstrom, K. Increased expression of endogenous biotin, but not BDNF, in telencephalic song regions during zebra finch vocal learning (120) 113

Joseph-Bravo, P., see Niquet, J. (120) 49 Juchau, M.R., see Trofimova-Griffin, M.E. (120) 7 Jungalwala, F.B., see Zhao, Z. (120) 165 Juraska, J.M., see Nuñez, J.L. (120) 87

Kawamura, K., see Fukuda, T. (120) 65
Kawano, H., see Fukuda, T. (120) 65
Ke, M., see Boylan, C.B. (120) 91
Kesterson, K.L., see Boylan, C.B. (120) 91
Kim, J.H.Y., see Nuñez, J.L. (120) 87
Kivell, B.M., McDonald, F.J. and Miller, J.H. Serum-free culture of rat post-natal and fetal brainstem neurons (120) 199
Kondo, H., see Saito, S. (120) 41

Lephart, E.D., see Lund, T.D. (120) 261
Levine, R.A., see Groc, L. (120) 95
Lindsley, T.A., see Yanni, P.A. (120) 233
Liu, Q.-Y., Coulombe, M., Dumm, J., Shaffer, K.M., Schaffner, A.E., Barker, J.L., Pancrazio, J.J., Stenger, D.A. and Ma, W.
Synaptic connectivity in hippocampal neuronal networks cultured on micropatterned surfaces (120) 223

Llamosas, M.M., Huerta, J.J., Cernuda-Cernuda,
R. and García-Fernández, J.M.
Ontogeny of a photic response in the retina and suprachiasmatic nucleus in the mouse
(120) 1

López-Costa, J.J., see Ríos, H. (120) 17
Lund, T.D., Salyer, D.L., Fleming, D.E. and Lephart, E.D.
Pre- or postnatal testosterone and flutamide effects on sexually dimorphic nuclei of the rat hypothalamus (120) 261

Ma, W., see Liu, Q.-Y. (120) 223

Manzanares, J., see Pérez-Rosado, A. (120) 77

McDonald, F.J., see Kivell, B.M. (120) 199

McEwen, B.S., see Orikasa, C. (120) 245

Miller, J.H., see Kivell, B.M. (120) 199

Mize, R.R., see Wu, H.H. (120) 105

Mooney, R.D., see Boylan, C.B. (120) 91

Moore, A.H., Hovda, D.A., Cherry, S.R., Villablanca, J.P., Pollack, D.B. and Phelps, M.E.

Dynamic changes in cerebral glucose metabolism in conscious infant monkeys during the first year of life as measured by positron emission tomography (120) 141

Muller, Y.L., Reitstetter, R. and Yool, A.J. Antisense knockdown of calcium-dependent K⁺ channels in developing cerebellar Purkinje neurons (120) 135 Nair, S.M., see Zhao, Z. (120) 165 Nakada, N., see Nishinaka, N. (120) 57 Nelson, J., see Nuñez, J.L. (120) 87 Niquet, J., Pérez-Martínez, L., Guerra, M., Grouselle, D., Joseph-Bravo, P. and Charli,

J.-L. Extracellular matrix proteins increase the

expression of pro-TRH and pro-protein convertase PC1 in fetal hypothalamic neurons in vitro (120) 49

Nishinaka, N., Hongo, S., Zhou, C.J., Shioda, S., Takahashi, R., Yamauchi, Y., Ohashi, T., Ohki, T., Nakada, N., Takeda, F. and Takeda, M.

Identification of the novel developmentally regulated gene, *Bdm2*, which is highly expressed in fetal rat brain (120) 57

Normile, H.J., see Groc, L. (120) 95 Norstrom, E., see Johnson, F. (120) 113 Nuñez, J.L., Nelson, J., Pych, J.C., Kim, J.H.Y.

and Juraska, J.M.

Myelination in the splenium of the corpus callosum in adult male and female rats (120)

Ohashi, T., see Nishinaka, N. (120) 57
Ohki, T., see Nishinaka, N. (120) 57
Orikasa, C., McEwen, B.S., Hayashi, H.,
Sakuma, Y. and Hayashi, S.
Estrogen receptor alpha, but not beta, is
expressed in the interneurons of the hippocampus in prepubertal rats: an in situ
hybridization study (120) 245
Osumi, N., see Fukuda, T. (120) 65

Pancrazio, J.J., see Liu, Q.-Y. (120) 223
Paquin, J., see Cadet, N. (120) 211
Pérez-Martínez, L., see Niquet, J. (120) 49
Pérez-Rosado, A., Manzanares, J., Fernández-Ruiz, J. and Ramos, J.A.
Prenatal Δ⁹-tetrahydrocannabinol exposure modifies proenkephalin gene expression in the fetal rat brain: sex-dependent differences (120) 77

Phelps, M.E., see Moore, A.H. (120) 141 Pollack, D.B., see Moore, A.H. (120) 141 Pomata, P.E., Colman-Lerner, A.A., Barañao, J.L. and Fiszman, M.L. In vivo evidences of early neurosteroid synthesis in the developing rat central nervous system and placenta (120) 83

Proctor, M.D., see Jaworski, D.M. (120) 27 Pych, J.C., see Nuñez, J.L. (120) 87

Rabey, J.M., see Gilad, V.H. (120) 255
Ramos, J.A., see Pérez-Rosado, A. (120) 77
Reitstetter, R., see Muller, Y.L. (120) 135
Rhoades, R.W., see Boylan, C.B. (120) 91
Ríos, H., López-Costa, J.J., Fosser, N.S., Brusco, A. and Saavedra, J.P.
Development of nitric oxide neurons in the chick embryo retina (120) 17
Rios, O., see Villalobos, J. (120) 99
Rizk, T., see Boylan, C.B. (120) 91

Saavedra, J.P., see Ríos, H. (120) 17 Saito, S., Sakagami, H. and Kondo, H. Localization of mRNAs for phospholipase D (PLD) type 1 and 2 in the brain of developing and mature rat (120) 41 Sakagami, H., see Saito, S. (120) 41 Sakuma, Y., see Orikasa, C. (120) 245 Salyer, D.L., see Lund, T.D. (120) 261 Schaffner, A.E., see Liu, Q.-Y. (120) 223 Seidler, F.J., see Zeiders, J.L. (120) 125 Shaffer, K.M., see Liu, Q.-Y. (120) 223 Shioda, S., see Nishinaka, N. (120) 57 Shuman, D.L., see Wu, H.H. (120) 105 Slotkin, T.A., see Zeiders, J.L. (120) 125 Soderstrom, K., see Johnson, F. (120) 113 Stenger, D.A., see Liu, Q.-Y. (120) 223 Strohl, K.P., see Thomas, A.J. (120) 181

Takahashi, R., see Nishinaka, N. (120) 57
Takeda, F., see Nishinaka, N. (120) 57
Takeda, M., see Nishinaka, N. (120) 57
Thomas, A.J., Erokwu, B.O., Yamamoto, B.K.,
Ernsberger, P., Bishara, O. and Strohl, K.P.
Alterations in respiratory behavior, brain
neurochemistry and receptor density induced by pharmacologic suppression of
sleep in the neonatal period (120) 181

Tobet, S., see Zhao, Z. (120) 165
Trofimova-Griffin, M.E., Brzezinski, M.R. and Juchau, M.R.
Patterns of CYP26 expression in human prenatal cephalic and hepatic tissues indicate an important role during early brain development (120) 7

Turner, J.E., see Jaynes, C.D. (120) 267

Villablanca, J.P., see Moore, A.H. (120) 141 Villalobos, J., Rios, O. and Barbosa, M. Postnatal development of the basal forebrain cholinergic projections to the medial prefrontal cortex in mice (120) 99

Weissmann, D., see Groc, L. (120) 95
Wiedenmayer, C.P., Goodwin, G.A. and Barr, G.A.
The effect of periaqueductal gray lesions on responses to age-specific threats in infant rats (120) 191

Wu, H.H., Cork, R.J., Huang, P.L., Shuman, D.L. and Mize, R.R. Refinement of the ipsilateral retinocollicular projection is disrupted in double endothelial and neuronal nitric oxide synthase gene knockout mice (120) 105

Yamamoto, B.K., see Thomas, A.J. (120) 181 Yamauchi, Y., see Nishinaka, N. (120) 57 Yanni, P.A. and Lindsley, T.A. Ethanol inhibits development of dendrites and synapses in rat hippocampal pyramidal neuron cultures (120) 233 Yool, A.J., see Muller, Y.L. (120) 135

Zeiders, J.L., Seidler, F.J. and Slotkin, T.A. Ontogeny of G-protein expression: control by β-adrenoceptors (120) 125

Zhao, Z., Chou, D.K.H., Nair, S.M., Tobet, S. and Jungalwala, F.B.
Expression of sulfoglucuronyl (HNK-1) carbohydrate and its binding protein (SBP-1) in developing rat cerebellum (120) 165
Zhou, C.J., see Nishinaka, N. (120) 57